| Mesa County Trend Analysis, 2017 Progress Toward Healthy People 2020 Targets or Colorado 2020 Targets | | | | | | | | | | | |
|--|---------|------------------------------------|-------------------|-------|-------|-------|-------|--------------------|-----------------------------|-----------------------------|-----------------------------------|
| Meternel and Child Health Indicator (Carted by 2017 Bealting | | Healthy People 2020 or Colorado | MCH Data By Year* | | | | | | Short Term Trend: | Long Term Trend: | 2017 Position Relative to 2020 |
| Maternal and Child Health Indicator (Sorted by 2017 Position Relative to 2020 Target) | of Data | 2020 Target | 2007 | 2013 | 2014 | 2015 | 2016 | 2017 | 2013 to 2017 | Percent Change 2007 to 2017 | Target |
| Perinatal mortality rate | 5 | 5.9 per 1,000 | 7.2 | 5.2 | 5.1 | 4.9 | 4.8 | 4.9 | Improved | -31.9% | Meets target |
| % births very low birth weight | 3 | 1.4% | 1.0% | 1.1% | 1.5% | 1.6% | 1.4% | 1.3% | Mixed | +30.0% | Meets target |
| % births preterm | 3 | 8.0% ^a | 8.8% | 8.0% | 8.1% | 8.5% | 8.2% | 7.5% | Mixed | -14.8% | Meets target |
| % infants placed to sleep on their backs | 2 | 75.8% | N/A | 83% | 88% | 91% | 87% | N/A ^b | Mixed ^c | N/A | Meets target ^d |
| % very low birth weight births in Level IIIs | 5 | 83.7% | 97.8% | 96.3% | 94.9% | 95.2% | 95.4% | 96.9% | No real change | -0.9% | Meets target |
| % children ages 5-14 overweight or obese | 3 | 25.0% ^a | N/A | NA | N/A | 22% | 21% | 22% | No real change ^e | N/A | Meets target |
| % infants who were ever breastfed | 2 | 84.5% ^a | 87% | 94% | 87% | 89% | 91% | N/A ^b | No real change ^c | +4.6% ^f | Meets target ^d |
| % cesarean deliveries among low-risk first births | 3 | 18.5% ^a | N/A | 13.9% | 13.6% | 13.7% | 14.8% | 14.6% | Worse | N/A | Meets target |
| Neonatal mortality rate | 5 | 4.1 per 1,000 | 4.2 | 2.3 | 2.3 | 2.1 | 2.8 | 2.9 | Worse | -31.0% | Meets target |
| Infant mortality rate | 5 | 6.0 per 1,000 | 5.8 | 4.3 | 4.2 | 4.0 | 4.6 | 5.0 | Worse | -13.8% | Meets target |
| Teen suicide death rate ages 15-19 | 5 | 10.2 per 100,000 ^g | 13.9 | DS | DS | DS | DS | 8.3 | Cannot ascertain | -40.3% | Meets target |
| Postneonatal mortality rate | 5 | 2.0 per 1,000 | 1.6 | 2.0 | 2.0 | 1.9 | 1.7 | 2.1 | Mixed | +31.3% | Close to target |
| % mothers discussed what to do if depressed during/after pregnancy | 2 | 85.8% ^a | N/A | 72% | 79% | 83% | 78% | N/A ^b | Mixed ^c | N/A | Close to target ^d |
| % mothers with appropriate weight gain during pregnancy | 1 | 38.0% ^a | N/A | 34.6% | 31.8% | 31.2% | 30.8% | 33.4% | Worse | N/A | Close to target |
| % children ages 1-14 with a medical home | 3 | 63.3% | N/A | N/A | 77% | 78% | 66% | 61% | Worse ^h | N/A | Close to target |
| % births low birth weight | 3 | 7.8% | 8.0% | 7.1% | 7.2% | 7.8% | 8.0% | 8.0% | Worse | +0.0% | Close to target |
| Teen birth rate ages 15-17 | 3 | 10.0 per 1,000 ^a | 27.0 | 18.6 | 17.6 | 16.7 | 14.5 | 13.0 | Improved | -51.9% | At some distance |
| % mothers who smoked 3 months before pregnancy | 1 | 12.2% | N/A | 18.0% | 19.2% | 20.8% | 19.5% | 17.6% | Mixed | N/A | At some distance |
| Injury hospitalization rate ages 10-19 | 3 | 226.9 ^a | 511.1 | 321.7 | 320.5 | 356.7 | 361.1 | 331.1 ⁱ | Worse | -35.2% | At some distance |
| % mothers overweight or obese before pregnancy | 1 | 38.0% ^a | N/A | 45.1% | 48.1% | 49.5% | 47.2% | 49.2% | Worse | N/A | At some distance |
| % teens who were bullied in past year | 1 | 22.0% ^a | NA | N/A | N/A | N/A | 31% | 27% | Cannot ascertain | N/A | At some distance |
| % mothers who smoked during pregnancy | 1 | 6.1% ^a | N/A | 14.2% | 15.7% | 16.1% | 15.4% | 13.0% | Mixed | N/A | Far from target |
| Teen motor vehicle death rate ages 15-19 | 3 | 12.4 per 100,000 ^g | 22.9 | 19.6 | 13.4 | DS | 10.5 | DS | Cannot ascertain | N/A | Unknown |
| Motor vehicle death rate ages 0-14 | 3 | 12.4 per 100,000 ^g | DS | 3.5 | DS | DS | 0.0 | DS | Cannot ascertain | N/A | Unknown |
| % children ages 6-11 physically active 60 min per day | 3 | 37.0% ^a | N/A | N/A | N/A | DS | DS | DS | Cannot ascertain | N/A | Unknown |
| % children ages 1-5 w/ parents asked by provider to fill out child q'naire | 3 | 52.0% ^a | N/A | N/A | DS | DS | DS | DS | Cannot ascertain | N/A | Unknown |

^a Colorado 2020 target.

^b 2015 PRAMS data were not available at time of trend analysis production.

 $^{\rm c}$ Short term trend is measured from 2013 to 2016.

^d Position relative to 2020 target is based on the data reported in 2016, not 2017.

^e Short term trend is measured from 2015 to 2017.

 $^{\rm f}$ Long term trend is measured from 2007 to 2016.

⁹ Healthy People targets are for all ages and are not specific to the age group shown.

^h Short term trend is measured from 2014 to 2017.

ⁱ Hospitalization data transitioned from ICD 9 to ICD 10 codes in October 2015; interpret 2017 data with caution.

DS denotes data that has been supressed due to small sample size. Supression criteria is unique to each data source.

N/A denotes data not available or applicable.

Cells shaded in light blue represent survey data results which require greater change over time

to be considered "Improved" or "Worse" compared to non-survey data.

* Refer to the accompanying data years table to determine the exact years for each value.

Additional Notes

Maternal and Child Health Indicator

Data are shown for indicators that have been included in the MCH County Data Sets for a number of years, or for new national or state MCH performance measures. Eight indicators are based on survey data available from the Pregnancy Risk Assessment Monitoring System, the Colorado Child Health Survey, and the Healthy Kids Colorado Survey: ever breastfed, infants placed to sleep on their backs, pregnancy depression, child development questionnaire, medical home, overweight or obese children, physically active children, and bullied teens. Data and short term trends for these indicators are lightly shaded. Most other indicators come from birth and death data, but some indicators (injury hospitalization) are derived from special registries or data sets. Refer to the *Definitions for Data in 2017 County Trend Analysis* handout for futher explanation of each indicator.

Number of Years of Data

Indicators use one, two, three, or five years of data. For example, the 2017 teen motor vehicle death rate for ages 15-19 is based on three years of data (2013, 2014, and 2015).

Healthy People 2020 or Colorado 2020 Target

Every ten years, Healthy People sets targets for numerous public health objectives. For 2010 there were 467 separate objectives; for 2020, the number increased to nearly 600. While many 2010 objectives were retained for 2020, the targets for nearly all of them have changed, and the new targets are virtually all more modest than those set for 2010. For more information, visit: http://www.healthypeople.gov/2020/default.aspx. Colorado 2020 targets are used in place of Healthy People 2020 targets where available.

MCH Data by Year

Each year when a county trend analysis is released the most recent available data are presented. The years of data that are available differ among indicators. For example, the most recent data available for the teen motor vehicle death rate for ages 15-19 is from 2015. Because this measure is a three-year rate, data from 2013, 2014, and 2015 are presented under the 2017 heading. As a general rule, the values shown beneath a column heading reflect data from at least two years earlier. Refer to the accompanying data years table to determine the exact years for each value.

Short Term Trend: 2013 to 2017

Short term trends are categorized as "Improved," "Worse," "Mixed," or "No real change." For a variety of reasons, some trends cannot be determined and are shown as "Cannot ascertain." The labeling of the change between 2013 and 2017 (or other years as footnoted) may be somewhat arbitrary. Values based on survey data need relatively large changes to obtain "Improved" or "Worse" labels because percentages shown are estimates that have confidence intervals. Values based on full count data (births, deaths, registries) require smaller changes to obtain labels denoting change. The 2013 and 2017 data values usually contain data from the previous 5 years, 3 years, or 1 year, so the trends described reflect changes in the data that are reported in the MCH Trend Analysis for 2013 and 2017, and do not represent changes that actually occurred between 2013 and 2017.

Long Term Trend: Percent Change 2007 to 2017

Long term trends are presented when data are available and comparable between 2007 and 2017. The percent change is the difference between the old and new estimates divided by the old estimate. The result is multiplied by 100 to yield a percent. The percent change is how much the measure has increased (+%) or decreased (-%) over the ten-year period. The 2007 and 2017 data values usually contain data from the previous 5 years, 3 years, or 1 year (children overweight or obese and infants placed to sleep on their backs contain data from 2 years), so the trends described reflect changes in the data that are reported in the MCH County Data Sets for 2007 and in the MCH Trend Analysis for 2017 and do not represent changes that actually occurred between 2007 and 2017.

2017 Position Relative to 2020 Target

"Meets target" indicates that the value shown in the MCH Trend Analysis released in 2017 is the same or better than the target. "Close to target" indicates that the value is within 20 percent of the target. "At some distance" describes values that are more than 20 percent away from the target, but are less than double the target for indicators requiring a decrease and are up to half the target for indicators requiring an increase. "Far from target" indicates a value that is either double or half the HP 2020 target, depending on whether the indicator requires a decrease or an increase. The values shown in 2017 are compared to the HP 2020 targets or the Colorado 2020 targets; see footnotes.

This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number H18MC00006, State Systems Development Initiative (SSDI). This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.

| Mesa County Trend Analysis, 2017 Data Years | | | | | | | | | | | |
|--|-----------------------|------------------|-----------|-----------|-----------|-----------|-----------|--|--|--|--|
| | Number of Years of | MCH Data By Year | | | | | | | | | |
| Maternal and Child Health Indicator | Data | 2007 | 2013 | 2014 | 2015 | 2016 | 2017 | | | | |
| Perinatal mortality rate | 5 | 2001-2005 | 2007-2011 | 2008-2012 | 2009-2013 | 2010-2014 | 2011-2015 | | | | |
| % births very low birth weight | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % births preterm | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % infants placed to sleep on their backs | 2 | N/A | 2010-2011 | 2011-2012 | 2012-2013 | 2013-2014 | N/A | | | | |
| % very low birth weight births in Level IIIs | 5 | 2001-2005 | 2007-2011 | 2008-2012 | 2009-2013 | 2010-2014 | 2011-2015 | | | | |
| % children ages 5-14 overweight or obese | 3 | N/A | N/A | N/A | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % infants who were ever breastfed | 2 | 2000-2004 | 2010-2011 | 2011-2012 | 2012-2013 | 2013-2014 | N/A | | | | |
| % cesarean deliveries among low-risk first births | 3 | N/A | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| Neonatal mortality rate | 5 | 2001-2005 | 2007-2011 | 2008-2012 | 2009-2013 | 2010-2014 | 2011-2015 | | | | |
| Infant mortality rate | 5 | 2001-2005 | 2007-2011 | 2008-2012 | 2009-2013 | 2010-2014 | 2011-2015 | | | | |
| Teen suicide death rate ages 15-19 | 5 | 2001-2005 | 2007-2011 | 2008-2012 | 2009-2013 | 2010-2014 | 2011-2015 | | | | |
| Postneonatal mortality rate | 5 | 2001-2005 | 2007-2011 | 2008-2012 | 2009-2013 | 2010-2014 | 2011-2015 | | | | |
| % mothers discussed what to do if depressed during/after pregnancy | 2 | N/A | 2010-2011 | 2011-2012 | 2012-2013 | 2013-2014 | N/A | | | | |
| % mothers with appropriate weight gain during pregnancy | 1 | N/A | 2011 | 2012 | 2013 | 2014 | 2015 | | | | |
| % children ages 1-14 with a medical home | 3 | N/A | N/A | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % births low birth weight | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| Teen birth rate ages 15-17 | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % mothers who smoked 3 months before pregnancy | 1 | N/A | 2011 | 2012 | 2013 | 2014 | 2015 | | | | |
| Injury hospitalization rate ages 10-19 | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % mothers overweight or obese before pregnancy | 1 | N/A | 2011 | 2012 | 2013 | 2014 | 2015 | | | | |
| % teens who were bullied in past year | 1 | N/A | N/A | N/A | N/A | 2013 | 2015 | | | | |
| % mothers who smoked during pregnancy | 1 | N/A | 2011 | 2012 | 2013 | 2014 | 2015 | | | | |
| Teen motor vehicle death rate ages 15-19 | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| Motor vehicle death rate ages 0-14 | 3 | 2003-2005 | 2009-2011 | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % children ages 6-11 physically active 60 min per day | 3 | N/A | N/A | N/A | 2011-2013 | 2012-2014 | 2013-2015 | | | | |
| % children ages 1-5 w/ parents asked by provider to fill out child q'naire | 3 | N/A | N/A | 2010-2012 | 2011-2013 | 2012-2014 | 2013-2015 | | | | |