

CHILD FATALITY PREVENTION SYSTEM: SUDDEN UNEXPECTED INFANT DEATH DATA, 2014 - 2018



COLORADO
Department of Public
Health & Environment

Introduction

The Child Fatality Prevention System (CFPS) is a statewide network that focuses on preventing child deaths. Housed at the Colorado Department of Public Health and Environment (CDPHE), CFPS consists of 43 local review teams, a 46-member State Review Team, and the CFPS state support team at CDPHE. Local teams include community members and field experts. These teams complete case reviews of infant, child, and youth deaths in Colorado to describe trends and patterns and create strategies to prevent future deaths. The CFPS State Review Team develops recommendations for the legislature on how to prevent child deaths in an annual legislative report.

The system reviews all deaths that occur in Colorado among infants, children, and youth under age 18. CFPS does not review deaths of Colorado residents that occur out of state. This is different from other reports of child death data and other Colorado government data sources. As a result, the data presented in this data brief might not match other statistics reported at both the state and national levels.

This data brief provides an overview of sudden unexpected infant death (SUID) data from CFPS. Additional CFPS data is available at:

www.cochildfatalityprevention.com/p/reports.html.

The impact of policies and systems on child deaths

Generations of social, economic, and environmental inequities contribute to some families losing infants, children, and youth.¹ When interpreting the data, it is critical to not lose sight of these systemic, avoidable, and unjust factors. These factors perpetuate the disparities observed in child deaths in Colorado. Researchers work towards understanding how geography, race, ethnicity, sexual orientation, and gender identity correlate with health. It is critical that data systems like CFPS identify, understand, and eliminate life-long inequities that persist across groups. When limitations in the data system exist due to how data is collected, or because data is not collected, CFPS strives to provide additional context and research about how inequities impact child deaths. By changing policies and systems that create and perpetuate inequities, CFPS can reduce the number of child deaths that occur in Colorado. Examples of these inequities include, but are not limited to:

RURAL AND FRONTIER GEOGRAPHY	RACE AND ETHNICITY	SEXUAL ORIENTATION AND GENDER IDENTITY
<p>Limited access to Level 1 trauma centers and mental and behavioral health services.²</p> <p>Increased stigma associated with mental illness and seeking help.³</p> <p>Longer response times by emergency medical services.⁴</p> <p>→ These and other factors contribute to higher death rates in rural areas, including suicide⁵ and passenger vehicle deaths.⁶</p>	<p>Historical trauma, racism, and discrimination.^{7,8}</p> <p>Limited access to high-quality education,⁹ employment opportunities,¹⁰ healthy foods,¹¹ culturally traditional foods,¹² and health care.¹³</p> <p>Chronic stress.¹⁴</p> <p>→ These factors result in lasting health impacts for people of color that include infant mortality,¹⁵ high rates of homicide and gun violence,¹⁶ and increased motor vehicle deaths.¹⁷</p>	<p>Discrimination, stigma, and bias.¹⁸</p> <p>Rejection from family, friends, and community.¹⁹</p> <p>Non-inclusive school curricula and anti-harassment policies.²⁰</p> <p>Insufficient access to LGBTQ+-informed health care.²¹</p> <p>→ This chronic social stress that LGBTQ+ children and youth experience influences health across the lifespan, including higher rates of suicide²² and substance use.²³</p>

Overview of SUID

Sudden unexpected infant death (SUID) describes deaths of infants under age 1 that occur suddenly and unexpectedly, whether explained or unexplained.²⁴ These deaths often occur during sleep, however this is not always the case. SUID is an umbrella term that combines deaths due to accidental suffocation and strangulation in bed (ASSB), sudden infant death syndrome (SIDS), and deaths due to undetermined causes.

What is SUID?

Deaths of infants under age 1 that occur suddenly and unexpectedly, whether explained or unexplained.



Accidental suffocation and strangulation in bed (ASSB): Suffocation by (1) soft bedding, pillow or waterbed mattress, (2) overlaying or rolling on top of or against infant while sleeping, or (3) wedging of an infant between two objects. ASSB also includes strangulation by asphyxiation.^{25,26}

Sudden infant death syndrome (SIDS): Assigned to deaths that cannot be explained after a thorough case investigation, including a death scene investigation, autopsy and review of the clinical history.²⁷ SIDS is sometimes known as “crib death” or “cot death.”

Undetermined causes: Assigned to infant deaths when the cause is unknown. This may occur when the requirements for a SIDS classification are not met.²⁸

SUID is a leading cause of infant mortality in the United States. Infant mortality indicates not just the health and wellbeing of families, caregivers, and their children, but is also considered an important indicator of overall health and wellbeing of a nation. Understanding why infants die from SUID in the United States and Colorado, among other leading causes of infant mortality, provide insights into ways in which we can improve health for all.

From 2014-2018, CFPS identified and reviewed 225 SUID. This represents 13.3% of all infant deaths (under age 1) in Colorado for the period. The annual rate of SUID occurring in Colorado among residents remained stable over this period (Table 1). The overall rate of SUID from 2014-2018 was 65.9 per 100,000 population. Colorado's rate of SUID is about one and a half-times less than the national SUID rate (100.5 per 100,000 population from 2013-2017).²⁹

Table 1. Rates of sudden unexpected infant death (SUID) occurring in Colorado among Colorado residents by year, 2014-2018.

Year of Death	n	Live Births	Rate*	95% Confidence Interval	
				Lower Limit	Upper Limit
2014-2018	215	326,256	65.9	57.1	74.7
2014	51	65,817	77.5	56.2	98.7
2015	35	66,567	52.6	35.2	70.0
2016	47	66,613	70.6	50.4	90.7
2017	43	64,388	66.8	46.8	86.7
2018	39	62,871	62.0	42.6	81.5

*Per 100,000 live births among residents in Colorado, 2014-2018.

Data sources: Colorado Child Fatality Prevention System and Vital Statistics Program, Colorado Department of Public Health and Environment.

SUID Risk and Protective Factors

In 2011, the American Academy of Pediatrics (AAP) developed recommendations for a safe infant sleeping environment to help reduce the risk of SUID. Recommendations include on-the-back sleep positioning, using a firm sleep surface, room-sharing without bed-sharing, and avoiding soft bedding and overheating.³⁰ With a quickly expanding body of SUID prevention research, the AAP expanded upon these recommendations in 2016.³¹ The 2016 A-level recommendations, listed below, were used for this report.

Level A Recommendations:

- Back to sleep for every sleep.
- Use a firm sleep surface.
- Room-sharing (to age 6 months) with the infant on a separate sleep surface (to 1 year).
- Keep soft objects and loose bedding away from the infant’s sleep area.
- Pregnant women should seek and obtain regular prenatal care.
- Avoid smoke exposure during pregnancy and after birth.
- Avoid alcohol and illicit drug use during pregnancy and after birth.
- Breastfeeding.
- Consider offering a pacifier at nap time and bedtime.
- Avoid overheating.
- Caregivers should immunize infants in accordance with AAP and CDC recommendations.
- Do not use home cardiorespiratory monitors as a strategy for reducing the risk of SIDS.
- Health care providers, staff in newborn nurseries and NICUs, and child care providers should endorse and model the SIDS risk-reduction recommendations from birth.

- Media and manufacturers should follow safe sleep guidelines in their messaging and advertising.
- Continue the “Safe to Sleep” campaign, focusing on ways to reduce the risk of all sleep-related infant deaths, including SIDS, suffocation, and other unintentional deaths. Pediatricians and other primary care providers should actively participate in this campaign.

Although the availability of sleep environment data varies by investigation, Table 2 indicates that none of the 225 infants who died between 2014 and 2018 slept in a space that met all of the AAP’s Level A Recommendations for a safe infant sleeping environment.

Table 2. Adherence to American Academy of Pediatrics 2016 Safe Infant Sleeping Environment Recommendations for SUID occurring in Colorado, 2014-2018.**

American Academy of Pediatrics 2016 Recommendation	Satisfied recommendation		Did not satisfy recommendation		Missing or unknown	
	n	Percent	n	Percent	n	Percent
All AAP recommendations satisfied	0	0.0	225	100.0	0	0.0
Infant and sleep environment recommendations						
Back to sleep for every sleep	120	53.3	56	24.9	34	15.1
Use a firm sleep surface	39	17.3	169	75.1	*	*
Room-sharing without bed-sharing is recommended	39	17.3	159	70.7	17	7.6
Keep soft objects and loose bedding out of the sleep environment	38	16.9	172	76.4	15	6.7
Consider offering a pacifier at nap time and bedtime	20	8.9	147	65.3	44	19.6
Caregiver-related recommendations						
Pregnant women should receive regular prenatal care (9 or more visits)	112	49.8	73	32.4	40	17.8
Breastfeeding is recommended	173	76.9	31	13.8	21	9.3
Avoid smoke exposure during pregnancy and after birth	78	34.7	88	39.1	59	26.2
Avoid alcohol or illicit drug use during pregnancy and after birth	107	47.6	78	34.7	40	17.8

*Data points with fewer than 3 observations are suppressed.

**Task force on Sudden Infant Death Syndrome (2016). *Pediatrics* 138(5)

Data source: Child Fatality Prevention System, Colorado Department of Public Health and Environment.

Demographic Characteristics

Age and Sex

Consistent with national trends, the majority of SUID occurred among those under five months of age (72.0%, n=162).³² Males account for the majority of SUID occurring in Colorado, representing 54.2% (n=122) of all SUID.


Race and Ethnicity

A note about terminology: Hispanic ethnicity as collected on the Colorado death certificate includes those that identify as Mexican, Mexican American, Chicano, Chicana, Puerto Rican, Dominican, Cuban, Central American, South American, Latin American, Spanish, and other Latin or Hispanic descent.³³ Additionally, “Latinx” and “Chicanx” are increasingly used gender inclusive terms, respecting those with a non-binary gender identity.^{34,35} To ensure clarity, this report uses “Hispanic” throughout the data section to reflect how CFPS data is collected from the death certificate and to align with terminology used in cited literature and research.³⁶

Between 2014 and 2018, the majority of infants who died by SUID in Colorado were non-Hispanic white (54.2%, n=122), 27.1% (n=61) were of Hispanic origin, 14.7% (n=33) were non-Hispanic Black, and 1.8% (n=4) were non-Hispanic American Indian or Alaska Native (AIAN).

Colorado observed a significant disparity in the rate of SUID by race and ethnicity (Figure 1). The rate of SUID among non-Hispanic Black infants was 3.3 times higher (191.9 per 100,000 live births) than for non-Hispanic white infants (58.5 per 100,000 live births). Additionally, the rate of SUID among AIAN infants was 2.9 times higher (169.3 per 100,000 live births) than for non-Hispanic white infants, although this difference was not statistically significant. However, these patterns are consistent with national data showing the highest SUID rates among AIAN and non-Hispanic Black infants.³⁷

Figure 1. Rates of SUID occurring in Colorado among Colorado residents by race and ethnicity, 2014-2018 (n=215)



*Error bars represent 95% confidence limits for rates.

Racial and ethnic disparities also exist for other leading causes of infant mortality, such as congenital malformations.³⁸ Researchers believe that factors that may contribute to these racial differences include low educational attainment, low income, inadequate prenatal care access and utilization, and paternal involvement in child-rearing.³⁹ However, studies examining these individual-level factors fail to fully explain the racial and ethnic differences in infant mortality.⁴⁰ Instead, research highlights the role that social determinants and contextual factors, particularly historical, community, and environmental inequities, play in infant mortality prevention.⁴¹

The increased likelihood of SUID among AIAN populations may stem from the historical loss of population, land, and culture that was endured by these communities.⁴² This historical trauma includes the forcible removal of indigenous AIAN people from their lands in the United States, and the forcible transfer of children from their families to boarding schools designed to strip them of their culture.⁴³ This trauma shapes the current societal context and leads to substantial socioeconomic and health disparities, including increased infant mortality.⁴⁴

People of color who self-report experiencing unfair treatment, racism, and discrimination across the lifespan have a greater risk for poor mental and physical health.⁴⁵ There is a growing body of research showing that the chronic stress created by these everyday experiences of racism contribute to allostatic load, or “wear and tear” on the body.⁴⁶ When a person is stressed, their body releases stress hormones, including epinephrine, adrenaline, and cortisol, which normally act to protect the body. However, experiencing chronic stress causes these hormones to continuously pump into the body, no longer protecting the individual, but rather causing damage.⁴⁷ These biological processes attributed to chronic stress are associated with negative birth outcomes, including low birthweight and preterm birth, which in turn influence the risk of SUID.⁴⁸

Increased cortisol levels due to chronic stress across the lifespan are further compounded by community level inequities, such as poverty. Black women have historically been disproportionately exposed to neighborhood poverty, a well-established risk factor for infant mortality.⁴⁹ This factor is completely independent of individual measures of socioeconomic status. Data show 18.7% of AIAN and 18.0% of Black Coloradans live below the poverty level, compared to 7.5% of non-Hispanic white Coloradans.⁵⁰

In addition to neighborhood poverty, racial residential segregation can determine infant mortality. It is largely driven by discriminatory federal, state, and local policies, such as redlining, that create unjust geographic divisions among racial and ethnic groups.^{51,52} Racial segregation leads to neighborhood disadvantage by concentrating neighborhood poverty, increasing exposure to environmental stressors such as air pollutants, creating barriers to and fewer opportunities for a healthy lifestyle, limiting access to health services, and increasing housing and food insecurity.⁵³ The consequences of residential segregation resulting from historical practices like redlining continue to reverberate throughout communities of color today. As just one example, the Colorado Behavioral Risk Factor Surveillance System (BRFSS) shows that 45.7% of AIAN and 36.2% of Black Coloradans are food insecure, meaning that they lack reliable access to affordable, nutritious food, compared to 19.2% of non-Hispanic white Coloradans.^{54,55} Such community inequities are associated with increased infant mortality among AIAN and Black populations.^{56,57}

Differences in the prevalence of on-the-back sleep positioning and other sleep environment factors across racial and ethnic populations may also contribute to disparities. Focus groups conducted with Black and AIAN families highlight that infant sleep decisions are often driven by culture and perceptions of what makes the infant comfortable and safe.⁵⁸

The Pregnancy Risk Assessment Monitoring System (PRAMS) tracks several aspects of infant sleep position and environments among families in Colorado. In 2018, 88.1% of mothers in Colorado reported that they most often lay their infants down to sleep on their backs, with 91.3% of non-Hispanic white mothers and 73.5% of Black mothers reporting on-the-back placement for sleeping. The regular use of a safe sleep location is more common among non-Hispanic white families. In 2018, 89.7% of mothers in Colorado reported usually placing

their infant to sleep in a crib, bassinet, or portable play yard in the last two weeks, with 93.1% of non-Hispanic white mothers and 85.9% of Black mothers reporting use of these sleep locations.⁵⁹ The use of soft bedding is also more common among Black families in Colorado, which is consistent with national trends.^{60,61} Too few AIAN mothers were surveyed by PRAMS to report in accordance with applicable privacy standards to share information about how their infants sleep. However, national research illuminates similar differences in safe sleep practices. For instance, AIAN mothers report the highest prevalence of bed sharing, when compared to other racial and ethnic groups.⁶² Based on this data, safe sleep education and awareness programs should collaborate with communities of color on the best methods to increase safe sleep practices.

Based on the research presented in this section, individual-level behaviors of parents and caregivers do not fully explain racial disparities in infant mortality. These racial disparities result from the social conditions facilitated by structural racism and discrimination.

Geography


Between 2014 and 2018, the majority of Colorado residents under age 1 who died by SUID in Colorado resided in an urban county (89.3%, n=192), while 8.4% (n=18) lived in a rural county, and 2.3% (n=5) lived in a frontier county. Although not statistically significant, the rate of SUID among infants living in a frontier county (78.4 per 100,000 live births) was higher than those living in a rural (59.9 per 100,000 live births) or urban county (66.3 per 100,000 live births). Readers should interpret this data with caution, as the frontier and rural rates represent very few deaths, decreasing the stability of these rates.

This rate data is consistent with national data showing higher SUID rates in rural areas, based on maternal residence as provided on the birth certificate.⁶³ Research indicates that geographic and social isolation and decreased access to health services experienced by rural communities contributes to higher rates of infant mortality.⁶⁴

SUID Investigative Circumstances

Figure 2 demonstrates the proportion of SUID occurring in Colorado by mechanism of death. Among the 225 SUID identified from 2014-2018, 44.9% (n=101) were attributed to ASSB. Of the remaining mechanisms, 42.7% (n=96) were attributed to undetermined causes and 8.4% (n=19) fell under the criteria for SIDS. The rate of SIDS has decreased since the early 1990s nationally and in Colorado, while the rates of SUID attributed to undetermined causes and ASSB have increased.⁶⁵ These changes are driven by improvements in investigations, a more thorough understanding of case definitions, and collecting more detailed information about safe sleep circumstances.

Figure 2. SUID occurring in Colorado by cause category, 2014-2018 (n=225)




CFPS has limited ability to determine the circumstances related to infant deaths when death scene investigators do not conduct a full investigation. Infant death scene investigations are critical to a comprehensive understanding of the circumstances and factors contributing to unexplained infant deaths. In 1996, the CDC developed the Sudden Unexplained Infant Death Investigation Reporting Form (SUIDIRF) (www.cdc.gov/sids/SUIDRF.htm) to aid in the investigation and understanding of SUID.⁶⁶

From 2014 to 2018, 94.2% (n=212) of all SUID in Colorado had a death scene investigation. The SUIDIRF was used in 49.3% (n=111) of the investigations and 31.6% (n=71) of investigations included a scene reenactment with a doll. Figure 3 demonstrates an increasing use of the SUIDIRF and doll reenactments as part of death scene investigations for SUID in Colorado. This suggests that scene investigators have an increased awareness of the importance of these tools in understanding the circumstances of these deaths.

The CFPS 2020 Legislative Report includes a recommendation to encourage and incentivize law enforcement agencies and coroner offices to use the SUIDIRF during infant death scene investigations. This will ensure that law enforcement officers and coroner investigators consistently collect circumstance data when investigating a suspected SUID.

Figure 3. Proportion of SUID occurring in Colorado by selected investigative methods and year, 2014-2018 (n=225)



Conclusion

SUID is a leading cause of infant mortality in the United States. From 2014 to 2018, SUID accounted for 13.3% of all infant deaths (under age 1) in Colorado. The highest SUID rates were observed among non-Hispanic Black and AIAN infants, children, and youth. The majority of SUID during this time period were due to accidental suffocation and strangulation in bed (ASSB) and undetermined causes. Additionally, none of the infants who died between 2014 and 2018 slept in a space that met all of the AAP’s Level A Recommendations for a safe infant sleeping environment. Upstream prevention strategies that address social and structural inequities can reduce SUID. To learn more about the prevention strategies recommended by CFPS, view the 2020 Legislative Report (www.cochildfatalityprevention.com/p/reports.html).

For more information and CFPS data, please contact the CFPS Support Team at the Colorado Department of Public Health and Environment:

Sasha Mintz, Child Fatality Prevention System Epidemiologist | sasha.mintz@state.co.us

References

1. Bailey, Z. D., Krieger, N., Agénor, M., Graves, J., Linos, N., & Bassett, M. T. (2017). Structural racism and health inequities in the U.S.A.: evidence and interventions. *The Lancet*, 389(10077), 1453-1463.
2. Rost, K., Fortney, J., Fischer, E., & Smith, J. (2002). Use, quality, and outcomes of care for mental health: The rural perspective. *Medical Care Research and Review*, 59(3), 231-265.
3. Cantrell, C., Valley-Gray, S., & Cash, R. E. (2012). Suicide in rural areas: risk factors and prevention. *Rural Mental Health: Issues, Policies, and Best Practices*. New York, NY: Springer.
4. Mell, H. K., Mumma, S. N., Hiestand, B., Carr, B. G., Holland, T., & Stopyra, J. (2017). Emergency medical services response times in rural, suburban, and urban areas. *JAMA surgery*, 152(10), 983-984.
5. Cantrell, C., Valley-Gray, S., & Cash, R. E. (2012). Suicide in rural areas: risk factors and prevention. *Rural Mental Health: Issues, Policies, and Best Practices*. New York, NY: Springer.
6. Beck, L. F., Downs, J., Stevens, M. R., & Sauber-Schatz, E. K. (2017). Rural and urban differences in passenger-vehicle-occupant deaths and seat belt use among adults—United States, 2014. *MMWR Surveillance Summaries*, 66(17), 1.
7. Palacios, J. F., & Portillo, C. J. (2009). Understanding Native women's health: Historical legacies. *Journal of Transcultural Nursing*, 20(1), 15-27.
8. Pager, D., & Shepherd, H. (2008). The Sociology of Discrimination: Racial Discrimination in Employment, Housing, Credit, and Consumer Markets. *Annual Review of Sociology*, 34, 181-209.
9. Williams, D. R., & Collins, C. (2016). Racial residential segregation: a fundamental cause of racial disparities in health. *Public Health Reports*, 116(5), 404-16.
10. Collins, C. A., & Williams, D. R. (1999, September). Segregation and mortality: the deadly effects of racism?. In *Sociological Forum* (Vol. 14, No. 3, pp. 495-523). Kluwer Academic Publishers-Plenum Publishers.
11. Larson, N. I., Story, M. T., & Nelson, M. C. (2009). Neighborhood environments: disparities in access to healthy foods in the U.S. *American journal of preventive medicine*, 36(1), 74-81.
12. Greder, K., de Slowing, F. R., & Doudna, K. (2012). Latina immigrant mothers: Negotiating new food environments to preserve cultural food practices and healthy child eating. *Family and Consumer Sciences Research Journal*, 41(2), 145-160.
13. White, K., Haas, J. S., & Williams, D. R. (2012). Elucidating the role of place in health care disparities: the example of racial/ethnic residential segregation. *Health Services Research*, 47(3pt2), 1278-1299.
14. Williams, D. R., & Mohammed, S. A. (2013). Racism and health I: Pathways and scientific evidence. *American behavioral scientist*, 57(8), 1152-1173.

-
15. Acevedo-Garcia, D., Lochner, K. A., Osypuk, T. L., & Subramanian, S. V. (2003). Future directions in residential segregation and health research: a multilevel approach. *American journal of public health*, 93(2), 215-221.
 16. Collins, C. A., & Williams, D. R. (1999, September). Segregation and mortality: the deadly effects of racism?. In *Sociological Forum* (Vol. 14, No. 3, pp. 495-523). Kluwer Academic Publishers-Plenum Publishers.
 17. King, M. (2017). Under The Hood: Revealing Patterns Of Motor Vehicle Fatalities In The United States. *Publicly Accessible Penn Dissertations*. 2396. Retrieved on June 19, 2019 from: repository.upenn.edu/edissertations/2396.
 18. Meyer, I. H. (2003). Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychological bulletin*, 129(5), 674.
 19. Kelleher, C. (2009). Minority stress and health: Implications for lesbian, gay, bisexual, transgender, and questioning (LGBTQ) young people. *Counselling psychology quarterly*, 22(4), 373-379.
 20. Sadowski, M. (2020). Safe is not enough: Better schools for LGBTQ students. *Harvard Education Press*.
 21. Kates, J., Ranji, U., Beamesderfer, A., Salganicoff, A., & Dawson, L. (2015). Health and access to care and coverage for Lesbian, Gay, Bisexual and Transgender (LGBT) individuals in the U.S.
 22. Hatzenbuehler, M. L., & Pachankis, J. E. (2016). Stigma and minority stress as social determinants of health among lesbian, gay, bisexual, and transgender youth: research evidence and clinical implications. *Pediatric Clinics*, 63(6), 985-997.
 23. Moazen-Zadeh, E., Karamouzian, M., Kia, H., Salway, T., Ferlatte, O., & Knight, R. (2019). A call for action on overdose among LGBTQ people in North America. *The Lancet Psychiatry*, 6(9), 725-726.
 24. Moon, R. Y., & Task Force on Sudden Infant Death Syndrome. (2016). SIDS and other sleep-related infant deaths: evidence base for 2016 updated recommendations for a safe infant sleeping environment. *Pediatrics*, 138(5), e2-e34.
 25. Moon, R. Y., & Task Force on Sudden Infant Death Syndrome. (2016). SIDS and other sleep-related infant deaths: evidence base for 2016 updated recommendations for a safe infant sleeping environment. *Pediatrics*, 138(5), e2-e34.
 26. Shapiro-Mendoza, C. K., Kimball, M., Tomashek, K. M., Anderson, R. N., & Blanding, S. (2009). U.S. infant mortality trends attributable to accidental suffocation and strangulation in bed from 1984 through 2004: are rates increasing?. *Pediatrics*, 123(2), 533-539.
 27. Willinger, M., James, L. S., & Catz, C. (1991). Defining the sudden infant death syndrome (SIDS): deliberations of an expert panel convened by the National Institute of Child Health and Human Development. *Pediatric Pathology*, 11(5), 677- 684.
 28. Schnitzer, P. G., Covington, T. M., & Dykstra, H. K. (2012). Sudden unexpected infant deaths: sleep environment and circumstances. *American Journal of Public Health*, 102(6), 1204-1212.

-
29. Centers for Disease Control and Prevention. Data and Statistics. (2019). Retrieved from www.cdc.gov/sids/data.htm.
 30. Task Force on Sudden Infant Death Syndrome. (2011). SIDS and other sleep-related infant deaths: expansion of recommendations for a safe infant sleeping environment. *Pediatrics*, 128, 1030-1039.
 31. Task Force on Sudden Infant Death Syndrome. (2016). SIDS and other sleep-related infant deaths: Updated 2016 recommendations for a safe infant sleeping environment. *Pediatrics*, 138(5), 1-12.
 32. Centers for Disease Control and Prevention. Data and Statistics. (2019). Retrieved from www.cdc.gov/sids/data.htm.
 33. Overview: Colorado Birth Certificates and Death Certificates, Vital Records. Retrieved from: drive.google.com/file/d/1GqFYL473YSJp-gwj9bE5arvNzoQg3O9K/view.
 34. Padilla, Y. (2016, April 18). What does “Latinx” mean? A look at the term that’s challenging gender norms. *Complex*. Retrieved from www.complex.com/life/2016/04/latinx.
 35. Salinas Jr, C., & Lozano, A. (2019). Mapping and recontextualizing the evolution of the term Latinx: An environmental scanning in higher education. *Journal of Latinos and Education*, 18(4), 302-315.
 36. Office of Health Equity, Colorado Department of Public Health and Environment, Health Inequities Fact Sheet 2019: Latinx Coloradans Fact Sheet. Retrieved from: drive.google.com/file/d/1z1b15A9hGaRvx4XTTa9BiPnz5lwjvfr/view.
 37. Parks, S. E., Lambert, A. B. E., & Shapiro-Mendoza, C. K. (2017). Racial and ethnic trends in sudden unexpected infant deaths: United States, 1995-2013. *Pediatrics*, 139(6).
 38. Mathews, T. J., MacDorman, M. F., & Thoma, M. E. (2015). Infant mortality statistics from the 2013 period linked birth/infant death data set. *National Vital Statistics Reports*, 64(9), 1-29.
 39. Matoba, N., & Collins Jr, J. W. (2017). Racial disparity in infant mortality. In *Seminars in Perinatology*, 41(6), 354-359. WB Saunders.
 40. Collins Jr, J. W., & David, R. J. (1990). The differential effect of traditional risk factors on infant birthweight among Blacks and whites in Chicago. *American Journal of Public Health*, 80(6), 679-681.
 41. Pickett, K. E., Collins Jr, J. W., Masi, C. M., & Wilkinson, R. G. (2005). The effects of racial density and income incongruity on pregnancy outcomes. *Social Science & Medicine*, 60(10), 2229-2238.
 42. Waletski, S. (2018). Historical Trauma in Native American Communities: Implications, Interventions, and Clinical Considerations. 2018 NCUR.
 43. Evans-Campbell, T. (2008). Historical trauma in American Indian/Native Alaska communities: A multilevel framework for exploring impacts on individuals, families, and communities. *Journal of interpersonal violence*, 23(3), 316-338.
 44. Campbell, C. D., & Evans-Campbell, T. (2011). Historical trauma and Native American child development and mental health: An overview. American Indian and Alaska Native children and mental health: Development, context, prevention, and treatment, 1-26.

-
45. Thomas, M. D., Michaels, E. K., Reeves, A. N., Okoye, U., Price, M. M., Hasson, R. E., ... & Allen, A. M. (2019). Differential associations between everyday versus institution-specific racial discrimination, self-reported health, and allostatic load among Black women: implications for clinical assessment and epidemiologic studies. *Annals of Epidemiology*, 35, 20-28.
 46. Williams, D. R., & Mohammed, S. A. (2013). Racism and health I: Pathways and scientific evidence. *American behavioral scientist*, 57(8), 1152-1173.
 47. Grimm, M., & Cornish, D. L. (2018). Infant Mortality and Racism in the United States. *International Journal of Undergraduate Research and Creative Activities*, 10(1), 2.
 48. Bartick, M., & Tomori, C. (2019). Sudden infant death and social justice: A syndemics approach. *Maternal & child nutrition*, 15(1), e12652.
 49. O'campo, P., Burke, J. G., Culhane, J., Elo, I. T., Eyster, J., Holzman, C., ...Laraia, B. A. (2007). Neighborhood deprivation and preterm birth among non-Hispanic Black and White women in eight geographic areas in the United States. *American Journal of Epidemiology*, 167(2), 155-163.
 50. U.S. Census Bureau; American Community Survey, 2018 American Community Survey 1-Year Estimates, Poverty Status in the Past 12 Months, Table S1701. Retrieved from data.census.gov.
 51. Polednak, A. P. (1996). Trends in U.S. urban Black infant mortality, by degree of residential segregation. *American Journal of Public Health*, 86(5), 723-726.
 52. Brown, K. S., Kijakazi, K., Runes, C., & Turner, M. A. (2019). *Confronting Structural Racism in Research and Policy Analysis*. Urban Institute. Retrieved from www.urban.org/sites/default/files/publication/99852/confronting_structural_racism_in_research_and_policy_analysis_0.pdf.
 53. Bailey, Z. D., Krieger, N., Agénor, M., Graves, J., Linos, N., & Bassett, M. T. (2017). Structural racism and health inequities in the U.S.A.: evidence and interventions. *The Lancet*, 389(10077), 1453-1463.
 54. Office of Health Equity, Colorado Department of Public Health and Environment, Health Inequities Fact Sheet 2019: American Indian & Alaska Native Coloradans Fact Sheet. Retrieved from drive.google.com/file/d/1unUSRd37yCMzGQY2Lhco7KzPDRoSMOri/view.
 55. Office of Health Equity, Colorado Department of Public Health and Environment, Health Inequities Fact Sheet 2019: Black/African American Coloradans Fact Sheet. Retrieved from drive.google.com/file/d/1s0dh56JHqKpbduwjh0_9MUVY_EfBysrg/view.
 56. Anthopolos, R., James, S. A., Gelfand, A. E., & Miranda, M. L. (2011). A spatial measure of neighborhood level racial isolation applied to low birthweight, preterm birth, and birthweight in North Carolina. *Spatial and Spatio-Temporal Epidemiology*, 2(4), 235-246.
 57. Holzman, C., Eyster, J., Kleyn, M., Messer, L. C., Kaufman, J. S., Laraia, B. A., ...Elo, I. T. (2009). Maternal weathering and risk of preterm delivery. *American Journal of Public Health*, 99(10), 1864-1871.

-
58. Herman, S., Adkins, M., & Moon, R. Y. (2015). Knowledge and beliefs of African-American and American Indian parents and supporters about infant safe sleep. *Journal of community health, 40*(1), 12-19.
 59. Pregnancy Risk Assessment Monitoring System Data: 2018. Center for Health and Environmental Data, Colorado Department of Public Health and Environment.
 60. Pregnancy Risk Assessment Monitoring System Data: 2018. Center for Health and Environmental Data, Colorado Department of Public Health and Environment.
 61. Shapiro-Mendoza, C. K., Colson, E. R., Willinger, M., Rybin, D. V., Camperlengo, L., & Corwin, M. J. (2015). Trends in infant bedding use: National Infant Sleep Position study, 1993-2010. *Pediatrics, 135*(1), 10.
 62. Bombard, J. M., Kortsmitt, K., Warner, L., Shapiro-Mendoza, C. K., Cox, S., Kroelinger, C. D., ... & Burley, K. (2018). Vital signs: Trends and disparities in infant safe sleep practices—United States, 2009-2015. *Morbidity and Mortality Weekly Report, 67*(1), 39.
 63. Ely, D. M., & Hoyert, D. L. (2018). Differences between rural and urban areas in mortality rates for the leading causes of infant death: United States, 2013-2015.
 64. Sparks, P. J., McLaughlin, D. K., & Stokes, C. S. (2009). Differential neonatal and postneonatal infant mortality rates across U.S. counties: the role of socioeconomic conditions and rurality. *The Journal of Rural Health, 25*(4), 332-341.
 65. Malloy, M. H., & MacDorman, M. (2005). Changes in the classification of sudden unexpected infant deaths: United States, 1992-2001. *Pediatrics, 115*(5), 1247-1253.
 66. Iyasu, S., Rowley, D. L., & Hanzlick, R. L. (1996). Guidelines for death scene investigation of sudden, unexplained infant deaths: recommendations of the Interagency Panel on Sudden Infant Death Syndrome. *Morbidity and Mortality Weekly Report: Recommendations and Reports, i*-22.