Rates* and Case Fatality Percents of Invasive *Haemophilus influenzae*by Age Group: Denver Metropolitan Area^ and Colorado, 2008

AGE GROUP (years)	Cases			Deaths		
	EIP^ No.	(EIP Rate*)	State No.	(State rate*)	State No.	(% of cases)
<=1	3	(4.1)	6	(4.2)	0	
2-4	3	(2.7)	3	(1.4)	0	
5-14	1	(0.3)	3	(0.4)	0	
15-24	2	(0.6)	4	(0.5)	0	
25-34	2	(0.6)	4	(0.6)	0	
35-49	2	(0.3)	3	(0.3)	1	(33.3)
50-64	10	(2.3)	15	(1.6)	1	(6.7)
65+	13	(6.0)	21	(4.3)	5	(23.8)
Total	36	(1.5)	59	(1.2)	7	(11.9)

^{*}Rates are per 100,000 population and based upon 2008-based population estimates, State Demography Office - DOLA ^EIP area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties.

Rates* of Invasive *Haemophilus influenzae* by County: Denver Metropolitan Area^, 2008

COUNTY OF RESIDENCE	Cases		
	No.	(Rate*)	
ADAMS	4	(0.9)	
ARAPAHOE	9	(1.6)	
DENVER	9	(1.5)	
DOUGLAS	5	(1.8)	
JEFFERSON	9	(1.7)	

^{*}Rates are per 100,000 population and based upon 2008-based population estimates, State Demography Office - DOLA ^EIP area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Sterile Sites from which *Haemophilus influenzae* was Isolated: Denver Metropolitan Area^, 2008

SITE OF ISOLATION	No.*	(% of cases)
Blood	32	(88.9)
CSF	3	(8.3)
Pleural fluid	3	(8.3)
Joint fluid	1	(2.8)
Placenta	1	(2.8)

*H.flu was isolated from one site in 30 cases and multiple sites in 6 case (Total cases=36)

H.flu was isolated only from blood in 26 cases (72.2%)

[^]Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Clinical Syndromes Associated with Invasive *Haemophilus influenzae*Disease:

Denver Metropolitan Area^, 2008

CLINICAL SYNDROME	No.*	(% of cases)
Bacteremia only	15	(41.7)
Bacteremic pneumonia	14	(38.9)
Meningitis	3	(8.3)
Cellulitis	1	(2.8)
Abscess	1	(2.8)
Pericarditis	1	(2.8)
Septic arthritis	1	(2.8)

*Some cases may have more than one syndrome (Total cases=36)
^Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Underlying Conditions Associated with Invasive *Haemophilus influenzae* :

Denver Metropolitan Area^, 2008

UNDERLYING CONDITION	No.*	(% of cases)
Diabetes	8	(22.2)
Current Smoker	6	(16.7)
Emphysema/COPD	6	(16.7)
Alcohol Abuse	4	(11.1)
Cerebral Vascular Accident/Stroke	4	(11.1)
Immunosuppressive Therapy	3	(8.3)
Asthma	3	(8.3)
Renal Failure/Dialysis	3	(8.3)
Obesity	3	(8.3)
Atherosclerotic Cardiovascular Disease	2	(5.6)
Heart Failure/CHF	2	(5.6)
Organ transplant	2	(5.6)
Cirrhosis/Liver Failure	1	(2.8)

*13 cases had more than one underlying condition
10 (27.8%) had no underlying conditions; 0 cases were unknown (Total cases=36)
^Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Invasive *Haemophilus influenzae* Serotypes: Denver Metropolitan Area^, 2008

Serotype	No.*	(% of isolates)
Not Typeable	23	(63.8)
f	10	(27.7)
а	2	(5.5)
b	1	(2.7)
Total	36	(100.0)

*Based upon testing of 36 isolates (Total cases=36)

Testing performed by the CDPHE Laboratory

^Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties