Rates* and Case Fatality Percents of Invasive *Haemophilus influenzae by Age Group:*

Denver Metropolitan Area^{and Colorado}, 2011

	Cases			Deaths		
AGE GROUP (years)	EIP^ No.	(EIP Rate*)	State No.	(State rate*)	State No.	(% of cases)
<=1	1	(1.5)	5	(3.8)	1	(20.0)
2-4					0	
5-14	2	(0.6)	2	(0.3)	0	
15-24	3	(1.0)	5	(0.7)	0	
25-34	3	(0.8)	4	(0.5)	0	
35-49	2	(0.4)	5	(0.5)	0	
50-64	8	(1.7)	19	(1.9)	4	(21.1)
65+	9	(3.5)	26	(4.5)	2	(7.7)
Total	28	(1.1)	66	(1.3)	7	(10.6)

*Rates are per 100,000 population and based upon 2011-based population estimates, State Demography Office - DOLA

^EIP area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties.

Rates* of Invasive *Haemophilus influenzae by County:*Denver Metropolitan Area^, 2011

COUNTY OF	Cases		
RESIDENCE	No.	(Rate*)	
ADAMS	5	(1.1)	
ARAPAHOE	6	(1.0)	
DENVER	10	(1.6)	
DOUGLAS	1	(0.3)	
JEFFERSON	6	(1.1)	

*Rates are per 100,000 population and based upon 2011-based population estimates, State Demography Office - DOLA

^EIP area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Sterile Sites from which *Haemophilus influenzae was Isolated:*Denver Metropolitan Area^, 2011

SITE OF ISOLATION	No.*	(% of cases)
Blood	26	(92.9)
CSF	1	(3.6)
Pleural fluid	1	(3.6)
Joint fluid	1	(3.6)

*H.flu was isolated from one site in 26 cases and multiple sites in 2 cases (Total cases=28)

H.flu was isolated only from blood in 24 cases (85.7%)

[^]Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Clinical Syndromes Associated with Invasive *Haemophilus influenzae Disease:*

Denver Metropolitan Area^, 2011

CLINICAL SYNDROME	No.*	(% of cases)
Bacteremic pneumonia	9	(32.1)
Bacteremia only	8	(28.6)
Meningitis	2	(7.1)
Epiglottitis	2	(7.1)
Septic abortion	2	(7.1)
Empyema	2	(7.1)
Otitis media	1	(3.6)
Peritonitis	1	(3.6)
Chorioamnionitis	1	(3.6)
Septic arthritis	1	(3.6)
Endocarditis	1	(3.6)

*Some cases may have more than one syndrome (Total cases=28)
^Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Underlying Conditions Associated with Invasive *Haemophilus influenzae :*Denver Metropolitan Area^, 2011

UNDERLYING CONDITION	No.*	(% of cases)
Current Smoker	7	(25.0)
Emphysema/COPD	6	(21.4)
Diabetes	3	(10.7)
Alcohol Abuse	3	(10.7)
Atherosclerotic Cardiovascular Disease	3	(10.7)
Cirrhosis/Liver Failure	2	(7.1)
Heart Failure/CHF	2	(7.1)
Cerebral Vascular Accident/Stroke	2	(7.1)
Hodgkin"s Disease	1	(3.6)
Asthma	1	(3.6)
Systemic Lupus Erythematosus	1	(3.6)
AIDS	1	(3.6)
Intravenous Drug Use	1	(3.6)
Obesity	1	(3.6)

*8 cases had more than one underlying condition 8 (28.6%) had no underlying conditions; 0 cases were unknown (Total cases=28) ^Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties

Invasive *Haemophilus influenzae Serotypes:*Denver Metropolitan Area^, 2011

Serotype	No.*	(% of isolates)
Not Typeable	15	(53.5)
f	9	(32.1)
а	2	(7.1)
е	2	(7.1)
Total	28	(100.0)

*Based upon testing of 28 isolates (Total cases=28)
Testing performed by the CDPHE Laboratory

^Emerging Infections Program area: Adams, Arapahoe, Denver, Douglas, and Jefferson counties