

Number of cases and rate of laboratory-confirmed and probable enteric infections, hospitalizations, and deaths, by disease – Communicable Disease Surveillance, Colorado 2017

| Disease | Cases | | | Hospitalizations | | Deaths | |
|---|--------------|-------|------------|------------------|--------|-----------|-------|
| | No. | Rate† | Objective§ | No. | (%) | No. | (%) |
| Bacteria | | | | | | | |
| <i>Botulism, Foodborne</i> | 0 | - | n/a | 0 | - | 0 | - |
| <i>Botulism, Infant</i> | 0 | - | n/a | 0 | - | 0 | - |
| <i>Campylobacter</i> | 1,092 | 19.5 | 8.5 | 177 | 16.2% | 2 | 0.2% |
| <i>Cholera</i> | 0 | - | n/a | 0 | - | 0 | - |
| <i>Listeria</i> | 10 | 0.2 | 0.2 | 10 | 100.0% | 2 | 20.0% |
| <i>Salmonella</i> | 686 | 12.2 | 11.4 | 153 | 22.3% | 5 | 0.7% |
| <i>Shigella</i> | 161 | 2.9 | n/a | 37 | 23.0% | 1 | 0.6% |
| Shiga toxin-producing <i>E. coli</i> (O157) | 60 | 1.1 | 0.6 | 28 | 46.7% | 1 | 1.7% |
| Shiga toxin-producing <i>E. coli</i> (non-O157) | 189 | 3.4 | n/a | 31 | 16.4% | 0 | 0.0% |
| <i>Typhoid Fever</i> | 5 | 0.1 | n/a | 5 | 100.0% | 0 | 0.0% |
| <i>Vibrio</i> | 37 | 0.7 | 0.2 | 2 | 5.4% | 0 | 0.0% |
| <i>Yersinia¶</i> | 13 | 0.4 | 0.3 | 3 | 23.1% | 0 | 0.0% |
| Parasites | | | | | | | |
| <i>Cryptosporidium</i> | 175 | 3.1 | n/a | 23 | 13.1% | 0 | 0.0% |
| <i>Cyclospora</i> | 27 | 0.5 | n/a | 1 | 0.37 | 0 | 0.0% |
| <i>Giardia</i> | 465 | 8.3 | n/a | 31 | 6.7% | 1 | 0.2% |
| Viruses | | | | | | | |
| <i>Hepatitis A</i> | 59 | 1.1 | n/a | 31 | 52.5% | 1 | 1.7% |
| Total | 2,979 | | n/a | 532 | | 13 | |

Abbreviations: N/A = not available

† Per 100,000 population.

§ Healthy People 2020 objective targets for incidence of *Campylobacter*, *Listeria*, *Salmonella*, STEC O157, *Vibrio*, and *Yersinia* infections per 100,000 population.

¶ *Yersinia* is only reportable in Adams, Arapahoe, Boulder, Broomfield, Denver, Douglas and Jefferson counties